EDITORIAL

What a year that was: some thoughts on health inequalities and publishing WEIRD articles

Theo Stickley

Joint Editor, Perspectives in Public Health

In this past year, global events have focused our minds on health inequalities. Specifically, the pandemic exposed the global inequalities surrounding access to vaccinations and healthcare; the energy, housing, and cost of living crises combined with soaring inflation in the West have further exposed national inequalities. Furthermore, the climate crisis focuses our minds on the fact that those who suffer the most from global heating (caused largely by the lifestyles and industrial activities of the wealthiest nations) are the poorest populations in the world. Social, economic and health inequalities are getting worse, not better.

The concept of health inequalities has been with us in the UK for decades. The 1980 Black report was the first report to clearly identify health inequalities in the modern age. The Acheson Report 18 years later found that the gap in inequalities in health had been steadily increasing and that differences in material deprivation are a major cause of the increase. Wilkinson¹ reflected on the continuous rise in social and health inequalities and astutely observed that political solutions were possible but not economically palatable to politicians. The Marmot² review was much more thorough than both the Black Report and the Acheson Report and presented findings and recommendations that were consistent across the 30-year period since the Acheson Report. In 2020, Marmot³ reviewed his findings and the progress made in the intervening decade. The findings are grim; poverty has risen and health inequalities worsened. The UK 'levelling-up' agenda falls a long way short of tackling inequalities.

Public Health has been defined as 'The art and science of preventing disease, prolonging life and promoting health through the organised efforts of society'.⁴ Is it time to redefine Public Health, or at least to include challenging and tackling inequalities? It is both poverty and inequalities that cause misery and hardship to billions of people on the planet, and the fight for survival is the fight for life itself.

Perspectives in Public Health is a journal that happily publishes research and practice articles that celebrate innovation in public health research and practice. But do we publish enough articles that acknowledge practice and research that identifies and tackles health inequalities? We openly acknowledge that our journal has a Western bias. It could be argued that much of what we publish is WEIRD (Western, Educated, Industrialised, Rich and Democratic according to Henrich et al.⁵). In a nutshell, this thesis is that researchers assume either there is little variation across populations or research subjects are universally representative. However, if researchers mainly study WEIRD populations, their findings will actually be skewed towards a minority population as the majority of the global population is not WEIRD; furthermore, as we always need to remind ourselves, non-White people are the global majority.

Sadly, the editorial team at *Perspectives* is constantly rejecting articles from around the world, not because we do not want to publish international research, but because often the quality of the research and reporting is not up to the standards we maintain. We wish to become less WEIRD and publish more research and practice that addresses inequalities and welcome submissions where research and practice participants are not WEIRD; we wish to contribute to the solution of inequalities and not become part of the problem.

For health professionals to tackle health and social inequalities, this inevitably involves political engagement. In the aftermath of the pandemic, the accelerating crisis of global heating, together with the current cost-of-living crisis, has accentuated the dilemmas health professionals frequently face. On the one hand, their commitment is to promote health, but on the other, they work in systems (micro, meso and macro) that maintain inequalities that, in turn, exacerbate ill-health within populations. Various current crises combine to create a public health emergency. In the light of increased concern over global public health, should health promotion of the future become more radical in order to help redress inequalities? With this in mind, this issue of *Perspectives in Public Health* begins with an opinion article from a practitioner who grapples with these issues in relation to climate change and offers food for thought for health practitioners around the world. Other topics in this issue include pre-natal exercises (Miguel Sánchez Polán et al), unintentional drowning in Spain (del Real et al), cervical cancer along the Texas-Mexico border (Salcedo et al) and the significance of friendships and mental health (Doran et al).

This year will undoubtedly bring further tests and challenges to public health around the world, and the editorial team here at *Perspectives* wish all our readers a happy new year.

REFERENCES

- 1. Wilkinson R. The impact of inequality: how to make sick societies healthier. New York: The New Press; 2005.
- 2. Marmot M. Fair society, healthy lives: the Marmot review: strategic review of health inequalities in England post-2010. Verona: Casa Editrice Leo S. Olschki s.r.l.; 2010.
- 3. Marmot M. Health equity in England: the Marmot review 10 years on. *BMJ* 2020;**368**:m693.
- 4. Acheson D. Public health in England: the report of the committee of inquiry into the future development of the public health function. London: The Stationary Office: 1988.
- 5. Henrich J, Heine SJ, Norenzayan A. The weirdest people in the world? Behav Brain Sci 2010;33(2-3):61-83.



The November 2022 CPD paper was 'The forensic implications of food hypersensitivity – a review of cases in United Kingdom courts: January 2014–February 2020' by MH Gowland et al.

Answers: 1c, 2a, 3d, 4d

Health professionals as activists: tackling threats to public health

In this short article, Katie Burns considers the role of healthcare professionals in light of the greatest global threat to public health: climate change.

K Burns

Teaching Associate, School of Health Sciences, The University of Nottingham, Queen's Medical Centre, Nottingham NG7 2RD, UK Email: katie.burns@nottingham.ac.uk

Corresponding author: *Katie Burns, as above*

The change in our climate is inescapable and affects the air we breathe, the water we drink, how much food we have to eat and whether we have safe places to live1 - these all being social and environmental determinants of health. Despite frequent warnings from meteorologists and other scientists, public attitudes and government policies in the northern hemisphere have been slow to change. Yet over the last few years, heat waves have been the cause of shortages of certain types of food,² and in the last year especially wildfires have destroyed homes, businesses and people's lives.³ Furthermore, the health of the public is in danger of deteriorating rapidly over the coming decades. It has recently been found that 58% of humanaffecting pathogenic diseases are exacerbated by climate-related changes.⁴ This is before we even begin to consider the psychological effects of

climate change and its effect on long-term conditions or the sourcing of treatments. The WHO predicts that before the first half of the century is over, the world will

The WHO predicts that before the first half of the century is over, the world will see 250,000 additional deaths per year due to malnutrition, vector-borne diseases and heat stress

see 250,000 additional deaths per year due to malnutrition, vectorborne diseases and heat stress.¹ But many more people, while not dying, will be living in poor health due to

changes in climate, and the majority of these will be among the poorest nations. With 90 scientific authors from 51 institutions around the world, the most recent *Lancet* countdown report presents data indicating these widespread changes in the environmental, social and economic determinants of health. The



Health professionals as activists: tackling threats to public health

report indicates the increasing impact climate change is having on both physical and mental health through the world's populations. These changes are often happening simultaneously, further accelerating exposure to health threats.⁵ What role is there for public health professionals in the face of such dire warnings?

A TIME FOR ACTION

Healthcare professions are among the most trusted individuals in society,⁶ and the time has come for significant radical change to what we consider as normal health promotion. It is therefore a moral imperative for healthcare workers to become more active in warning of the threats of global warming and to call governments to account for inaction. There hasn't previously been such a strong need for healthcare professionals to become activists to promote the health of the public and help prevent suffering and deaths in the future. A recent article in *Nature Climate Change*⁷ suggests that adding the voices of scientists and healthcare workers strengthens the argument for policy-makers to hear. The authors also advocate for civil

References

- World Health Organization. Climate change and health, 2021. Available online at: https:// www.who.int/news-room/fact-sheets/detail/ climate-change-and-health (Last accessed 9th September 2022).
- Horton H. UK farmers count cost as heatwave kills fruit and vegetable crops. *The Guardian* 2022; 1 August. Available online at: https:// www.theguardian.com/environment/2022/ aug/01/uk-farmers-count-cost-as-heatwavekills-fruit-and-vegetable-crops (Last accessed 9th September 2022).
- Gair K, Brown D, Parker C et al. UK heatwave: forty homes destroyed by wildfires in London. *The Times* 2022; 20 July. Available online at:

disobedience to strengthen the messaging. Groups such as Doctors for Extinction Rebellion cite the General Medical Council (GMC) code of conduct as their reason for speaking

out about the need for greater action from governments:

Our code of conduct compels us to act where we notice unacceptable risks to current and future patient health, and act promptly. We

have noted that traditional techniques of writing academic papers and journal articles have not produced sufficient meaningful results to continue with them alone.⁸

For those of us who genuinely care about global inequalities, the case for action is unequivocal as there is much evidence to indicate that those in the poorest countries will continue to suffer the most. This emerging fact was identified by the Intergovernmental Panel on Climate Change over 20 years ago.⁹ In the intervening period, little has

> https://www.thetimes.co.uk/article/ uk-heatwave-record-high-of-40-3c-sparkswildfires-in-tinderbox-britain-rs6z3crdc (Last accessed 9th September 2022).

- Mora C, McKenzie T, Gaw I et al. Over half of known human pathogenic diseases can be aggravated by climate change. Nat Clim Chang 2022;12:869–75.
- Romanello M, Napoli C, Drummond P et al. The 2022 report of the Lancet Countdown on health and climate change: health at the mercy of fossil fuels. The Lancet 2022. Available online at: https://www.thelancet.com/journals/ lancet/article/PIIS0140-6736(22)01540-9/ fulltext (Last accessed 8th November 2022).

changed for the plight of millions of people. Never before has there been such an urgent moment for healthcare workers to be involved in promoting health and preventing disease and

Never before has there been such an urgent moment for healthcare workers to be involved in promoting health and preventing disease and death, before it is too late death, before it is too late. In short, healthcare workers have a responsibility to become activists to help bring about lasting change, and the sooner, the better.

FUNDING

The author(s) received no financial support for the research, authorship and/or publication of this article.

CONFLICT OF INTEREST

The author(s) declared no potential conflicts of interest with respect to the research, authorship and/or publication of this article.

ORCID ID

K Burns (D) https://orcid.org/0000-0001-8312-1884

- Ipsos. Global Trustworthiness Index, 2021. Available online at: https://www.ipsos.com/ en-uk/doctors-become-worlds-most-trustedprofession (Last accessed 3rd October 2022).
- Capstick S, Thierry A, Cox E *et al.* Civil disobedience by scientists helps press for urgent climate action. *Nat Clim Chang* 2022;**12**:773–74.
- Doctors for Extinction Rebellion. Available online at: https://www.doctorsforxr.com/ (Last accessed 9th September 2022).
- Houghton JT, Ding Y, Griggs DJ et al. Climate change 2001: the scientific basis. Intergovernmental Panel on Climate Change, Cambridge, UK, January 2001.

IN PRACTICE

In Practice

Narrowing the gap between research and policy: using rapid evaluation during the COVID-19 crisis

K Anil

School of Health Professions, Faculty of Health and Human Sciences, University of Plymouth, Peninsula Allied Health Centre, Derriford Road, Plymouth PL6 8BH, UK Email: krithika anil@plymouth.ac.uk

Email: krithika.anil@plymouth.ac.uk

D Watson

Global Health Research Institute, School of Human Development and Health, Faculty of Medicine, University of Southampton, UK

J Alagil

School of Health Sciences, Faculty of Life and Environmental Sciences, University of Southampton, UK College of Applied Medical Sciences, Health Rehabilitation Department, King Saud University, Kingdom of Saudi Arabia

R Dewar-Haggart

School of Primary Care, Population Sciences and Medical Education, Faculty of Medicine, University of Southampton, UK

S Fearn

School of Human Development and Health, Faculty of Medicine, University of Southampton, UK

C McGrath

NIHR Southampton Biomedical Research Centre, University of Southampton and University Hospitals Southampton NHS Foundation Trust, UK MRC Lifecourse Epidemiology Unit, University of Southampton, UK

C Meagher

School of Health Sciences, Faculty of Life and Environmental Sciences, University of Southampton, UK

S Muir

MRC Lifecourse Epidemiology Unit, University of Southampton, UK

M Barker

School of Health Sciences, Faculty of Life and Environmental Sciences, University of Southampton, UK

NIHR Southampton Biomedical Research Centre, University of Southampton and University Hospitals Southampton NHS Foundation Trust, UK

MRC Lifecourse Epidemiology Unit, University of Southampton, UK

*D Watson is also affiliated to SAMRC Developmental Pathways for Health Research Unit, School of Clinical Medicine, University of the Witwatersrand, Johannesburg, South Africa and Department of Global Health and Social Medicine, King's College London, United Kingdom

Corresponding author:

Krithika Anil, as above

INTRODUCTION

An evidence base is vital to ensure well-informed policies with systematic processes.^{1,2} However, a research-to-policy gap exists that is widened by the discrepancy between the extensive time it takes to conduct conventional academic research and the short timescale over which policy-makers are often

required to make decisions.² This is especially concerning in times of crisis, such as the COVID-19 pandemic, that demanded an avalanche of data, analysis, and interpretation to be provided over a very short period of time. Rapid evaluation can be used to generate research-based evidence under pressure to inform decisionmaking and policy. Existing literature clearly details the steps involved in conducting rapid evaluation³; yet this literature contains little practical knowledge about how best to carry out such research. In this article, we reflect upon the practical implementation of rapid evaluation for an urgent project during the crisis of COVID-19.⁴

BRIEF PROJECT BACKGROUND AND REFLECTION PROCESS

Between June and October 2020, a COVID-19 Saliva Testing Programme was piloted, where findings were reported to the UK Department of Health and Social Care (DHSC).⁵ A rapid evaluation was conducted with participants of the testing programme to generate insights that would inform the testing's design and modification, and the next phase of future mass-testing to the UK DHSC.⁴ For further details on the project, see our published paper.⁴ We used Gibb's reflective cycle^{6,7} as a

A research-to-policy gap exists that is widened by the discrepancy between the extensive time it takes to conduct conventional academic research and the short timescale over which policy-makers are often required to make decisions foundation for the team to reflect on their experiences of working on a rapid qualitative project. The main 'lessons learned' are explained below, followed by actionable suggestions in Table 1.

CREATING AN EFFECTIVE TEAM FOR RAPID OUTPUTS

The team identified four main lessons critical to the success of our rapid evaluation project during a crisis: (1) the shared vision for the project held by the team, (2) effective project management, (3) the supportive leadership style, and

14 Perspectives in Public Health | January 2023 Vol 143 No 1



Table 1

Actionable points based on our reflection categories

Category	Specific actions for a leader	Specific actions for a team member
The Value of a Shared Vision	Clearly communicate the project's goals and its importance at the start and throughout the project to maintain team motivation	Ensure you know why the project is needed and be aware of project goals from the beginning
	 Plan the project in-depth as early as possible; this includes the following: Preparing a timeline for each step of the project Deciding deadlines Building the team and identifying individual member's strengths Deciding task delegation to the team based on skill 	Familiarise yourself with project timelines and deadlines
	 Create regular meeting spaces with team members to communicate the following: Project goals Deadlines Update on project outputs and next steps 	Ensure you know how to access project information and learn to use any required project-related technology
	Consider using a platform (such as Microsoft Teams) as a central location to store and share real-time project information and documents with the team	
	Following each meeting, send a summary and action points to all team members	
Supportive Leadership Style	Identify skill gaps among your team and provide support/training to reduce those gaps	Familiarise yourself with your team members' skills to identify opportunities to assist your own work and/or improve your own skills
	Listen to the opinions of your team and allow for the potential for their opinions to influence the project plans	Listen to the opinions of other team members and communicate challenges to facilitate collaborative problem solving
	Be prepared to listen to opposing opinions and make final decisions on any differences; this will be made easier by being clear on your project goals	
	Approach your senior colleagues for support and advice when needed	
	Recognise and show appreciation of team members for their efforts by providing positive feedback	Offer support to other team members where needed and if you have the capacity
	Reflect on your leadership skills, recognising your concerns and identifying ways to overcome them; use strategies such as Gibb's Reflective Cycle for reflection	 Noting the leadership skills of your project lead may help you vicariously develop your own leadership skills; look out for the following to help you: Methods used to communicate information Storage of information How they use technology to support the project How they interact with team members and the type of support they offer
Access to Rapid Evaluation Methods and Digital Technology	Understand the principles of a rapid approach and share them with the team at the beginning of the project to set expectations	Familiarise yourself with the principles of a rapid approach and be prepared to meet its expectations
	 Clearly communicate deadlines and expectations of work timescales to your team at the start; do this by: Storing deadlines and timelines on a central storage system that is easily accessible Ensuring every team member knows where and how to access this information Emphasising task expectations by outlining specific task details and workload involved Setting clear expectations of the work schedule and arranging working hours based on your team's availability 	Ensure you are prepared to reply to communications from your project lead and team members quickly to ensure task deadlines are met
	Balance your rapid project with your other work commitments by setting specific hours and days to work on specific projects	Communicate your specific working hours to your project lead to set communication expectations
	Update project information as soon as possible and communicate these updates to your team; using a real-time reporting system (e.g. on Microsoft Teams) will greatly assist this communication	
	Create a plan that can be flexible, based on your deadlines, your team's working hours, and skills of team members	

Figure 1



(4) access to rapid evaluation methods and digital technology.

The value of a shared vision

The pandemic crisis gave the work an urgency and significance felt by the team, resulting in both a personal and professional commitment to the project. Team members felt positive about the contribution they were making to the national campaign to reduce spread of the infection. This drive facilitated a proactive and supportive work culture, where team members responded to communications quickly and worked collectively to solve problems. All team members listened and valued opinions from other members and were quick to take up responsibilities if others did not have the capacity. A sense of trust and confidence were quickly built among the team members, who easily relied on each other to complete tasks and receive advice. The culture enabled the project to collect data beyond its target and produce outputs that were valued by DHSC for their timeliness and insights.

Effective project management

Effective project management and accessible information sharing and communication were key to generating project insights within the tight deadlines required by the stakeholders. The team leader communicated the project's goals and objectives at the beginning of the project, specifying how the project was to be completed. She also held weekly meetings to discuss experiences or challenges and real-time digital summaries accessible to all on Microsoft (MS) Teams. The well-organised communication of the project ensured all team members were up-to-date on the project and knew its immediate next steps. The team also had immediate access to all project-related documents and data via MS Teams, such as project protocols, task delegation spreadsheets, deadlines, and result summaries.

The supportive leadership style

The team leader initially identified the strengths, skills, backgrounds, and experience of team members in order to delegate tasks accordingly. The team leader's in-depth knowledge of her team led to inherent confidence in each member's ability to complete the project's tasks on time. The team consisted of 12 researchers from varied backgrounds and seniority that required active management from the team leader by listening to their opinions and concerns, and making members feel valued. Some team members were inexperienced with rapid evaluation methods and were initially worried about their contribution to the project. The team leader reassured them by stating

why they were asked to be on the team and provided appropriate support where needed. When team members had opposing perspectives on aspects of the project, the team leader made the final decision through team discussion – a process made easy due to the trust built over the course of the project.

Encouraging team members to actively participate and support each other eased the management burden of a large team for the team leader, which prevented her feeling overwhelmed with project speed and enabled her instead to enjoy the process. The team leader was open about gaps in her knowledge and sought help when needed. These leadership skills were respected by the team, who felt they vicariously learned about effective leadership and believed they would be employing these skills in future team management activities.

The relationship between the topics discussed in the above sections is shown in Figure 1.

Access to rapid evaluation methods and digital technology

Successfully implementing rapid evaluation requires access to the right resources that can be summarised as having a large team, using rapid assessment procedure ('RAP') sheets to analyse findings iteratively, and providing ongoing immediate feedback to stakeholders.⁸ Our project was entirely virtual due to the COVID-19 lockdown and therefore demanded reliance on technology. We used the synchronous features of MS Teams to share documents, communicate quickly, and keep track of tasks and deadlines. Our team were familiar with MS Teams but it is important to consider time to support any team members who are not familiar with the technology used for any project.

All team members were working remotely, some of whom had caring duties and other work responsibilities, which resulted in a flexible working timetable. Clear and strict deadlines were communicated in advance, and responsibilities were delegated based on skillset and availability as indicated on a live team calendar – a crucial resource for our project.

CHALLENGES CREATED BY THE RAPID EVALUATION PROCESS

Team members felt pressured to be on alert for communications and reply quickly, even outside of their normal working hours due to the flexible work schedule. Some data collection tasks were scheduled quickly, which was sometimes too abrupt for some members who could not make those timings. The team leader felt particular pressure as she had to juggle leading this project with other work responsibilities. Finally, this project was ethically approved as a service evaluation and not as research because service evaluation ethics was quicker to attain. This restricted dissemination of this project's findings and delayed publication. Despite these challenges, team members openly discussed and overcame these challenges because of the culture of trust and support built from the start of the project.

CONCLUSION

Our experience illustrates how a rapid yet rigorous evaluation to inform policy can generate actionable results under time pressure that directly informed government policy and practice. This

demonstration of how these outcomes can be achieved using entirely online methods suggests that a rapid approach to programme evaluation can be used irrespective of the location of research or public health partners, while also accommodating needs for flexible working. Furthermore, this approach makes the most of short contracts that govern much commissioned research in the UK. We have generated a list of actionable points based on our reflections on the conduct of this project which we hope could be used by anyone planning to conduct a rapid evaluation project to inform policy and practice regardless of the field of study (Table 1).

ACKNOWLEDGEMENTS

We would like to sincerely thank all our team members from the original project

they were a pleasure to work with: Natalia Laverty Baralle, Sandy Ciccognani, Julia Groot, Kathryn Knowles, Jo Musgrove, Kate Glyn-Owen, Kath Woods-Townsend, Andrew Mortimore, Paul Roderick, Janis Baird, Hazel Inskip, and Keith Godfrey.

ORCID IDS

A rapid approach to

programme evaluation

can be used

irrespective of the

location of research or

public health partners,

while also

accommodating needs

for flexible working

Krithika Anil D https://orcid.org/0000-0002-8027-1665 Rachel Dewar-Haggart https://orcid .org/0000-0002-3757-1152

References

- Oxman AD, Lavis JN, Lewin S et al. SUPPORT Tools for evidence-informed health policymaking (STP) 1: what is evidenceinformed policymaking? *Health Res Policy Syst* 2009;7(1):S1.
- Uzochukwu B, Onwujekwe O, Mbachu C et al. The challenge of bridging the gap between researchers and policy makers: experiences of a Health Policy Research Group in engaging policy makers to support evidence informed policy making in Nigeria. *Globalization and Health* 2016;**12**(1):67.
- 3. Vindrola-Padros C. *Doing rapid qualitative research*. Thousand Oaks, CA: Sage; 2021.
- Watson D, Baralle NL, Alagil J et al. How do we engage people in testing for COVID-19? A rapid qualitative evaluation of a testing programme in schools, GP surgeries and a university. BMC Public Health 2022;22(1): 305.
- Godfrey K. Evaluation of the expanded Southampton pilot study (Phase 2) for use of saliva-based lamp testing in asymptomatic populations: final report, 16th November 2020. Available online at: https://eprints. soton.ac.uk/445622/1/Evaluation_of_the_ expanded_Southampton_pilot_study_ Phase_2_for_use_of_saliva_based_lamp_

testing_in_asymptomatic_populations_eprints .pdf

- Gibbs G. Learning by doing: a guide to teaching and learning methods. Oxford: Further Education Unit; 1988.
- University of Edinburgh. Gibbs' reflective cycle, 2020. Available online at: https://www.ed. ac.uk/reflection/reflectors-toolkit/reflecting-onexperience/gibbs-reflective-cycle (Last accessed 2021 October).
- Vindrola-Padros C, Chisnall G, Cooper S et al. Carrying out rapid qualitative research during a pandemic: emerging lessons from COVID-19. *Qual Health Res* 2020;**30**(14):2192–204.

Barriers and facilitators to delivering bystander cardiopulmonary resuscitation in deprived communities: a systematic review

Authors

I Uny២

Institute for Social Marketing and Health, Faculty of Health Sciences and Sport, University of Stirling, Stirling FK9 4LA, UK. Email: isabelle.uny@stir.ac.uk

K Angus¹⁰ Institute for Social Marketing and Health, Faculty of Health Sciences and Sport, University of Stirling, Stirling, UK

E Duncan

Nursing, Midwifery and Allied Health Professions Research Unit, Faculty of Health Sciences and Sport, University of Stirling, Stirling, UK

F Dobbie

Usher Institute, College of Medicine and Veterinary Medicine, University of Edinburgh, Edinburgh, UK

Corresponding author: *Isabelle Uny, as above*

Keywords

cardiopulmonary resuscitation; CPR; bystander CPR; health inequalities; health improvement; emergency

Abstract

Background: There is a higher incidence of cardiac arrest in economically deprived areas; however, data show that bystander cardiopulmonary resuscitation (CPR) in those areas is lower. This results in lower survival rates, placing those communities at a double disadvantage. This systematic review explored the barriers and facilitators to engaging with bystander CPR in deprived communities.

Methods: Studies were eligible for inclusion if they addressed any barrier or facilitator to performing bystander CPR or being trained in CPR or training others. Studies had to either be set in a deprived area or examine a deprived population. Selected studies were published between January 2000 and December 2017 and reported on primary research. No language limitations were applied. Searches were conducted in the following databases: Cumulative Index to Nursing and Allied Health Literature (CINAHL), MEDLINE, PsycINFO, PubMed, and Web of Science Core Collection. Unpublished 'grey' literature was also searched as well as the reference lists of any relevant studies.

Results: The systematic review highlighted several main factors acting as barriers or facilitators to engaging with bystander CPR in deprived communities: (1) the willingness to learn or perform CPR, (2) the confidence to perform CPR, and (3) self-reported likelihood of performing CPR. The review also revealed additional barriers to engaging with CPR which are specific to – or more acute for – individuals from socioeconomically deprived backgrounds or areas.

Discussion: We found little evidence suggesting that the willingness to perform or learn bystander CPR is lower in deprived communities compared to the general population. However, the confidence to perform CPR in deprived communities was affected by some measures of socioeconomic status. The results also crucially highlighted other barriers more acute in deprived communities: the risk to personal safety in administering CPR; the fear of legal consequences; and the lack of community cohesion and other cultural barriers.

INTRODUCTION

Out-of-hospital cardiac arrests (OHCAs) are a major cause of mortality worldwide, and the burden they represent on health services is not fully known.^{1,2} Globally, survival from OHCA remains low and variable within regions with estimates ranging from 7.6% in Europe, 6.8% in North America, 3.0% in Asia, and 9.7% in Australia.¹

OHCA survival rates remain poor and have seen little improvement in recent decades.³ One of the main factors influencing the rate of survival is the administration of bystander cardiopulmonary resuscitation (CPR).⁴ There is an apparent willingness among the general population to administer bystander CPR,⁵ born out of a desire to save lives especially if the victim

Copyright © Royal Society for Public Health 2022 SAGE Publications © © © ISSN 1757-9139 DOI: 10.1177/17579139211055497 January 2023 Vol 143 No 1 | Perspectives in Public Health 43

review was to identify the barriers and

facilitators perceived by individuals in

Therefore, the aim of this systematic

is a family member.⁶ However, a number of barriers often prevent people from performing bystander CPR, regardless of whether they are CPR trained or not. Some of these barriers include a reluctance to perform CPR on a stranger,⁵ particularly if the victim seems unkempt or under the influence of drugs or alcohol,⁷ a fear of being sued,^{8,9} or a fear of performing CPR wrongly and of potentially doing more harm than good.^{5,10,11}

Countries that have renewed their efforts to increase the number of people trained in CPR have seen significant improvements in their OHCA survival rates. For instance, in Sweden, a CPR training policy which saw a third of the total population trained over three decades resulted in an increase in OHCA survival at 1 month from 5% in 1992 to 11% in 2011.^{12,13} A similar strategy in Denmark resulted in a rise in overall survival to discharge from hospital from 6.5% to 19.1% between 2001 and 2010¹⁴.

Rates of bystander CPR are generally lower in socially and economically deprived areas, in rural areas, and within certain ethnic populations.^{9–11} Sasson et al.¹⁵ found that victims of OHCA in low-income black neighbourhoods in the USA were less likely to receive bystander CPR than those in high-income white neighbourhoods. Another study in North East England found that residents living in the least deprived areas were significantly more likely to receive bystander CPR than those in the most deprived.¹⁶

Besides the fact that they are less likely to receive CPR, people living in socially deprived communities are also less likely to be trained in CPR. Studies have found that lower education and lower incomes were associated with a decreased likelihood of being trained in CPR.^{17,18} Furthermore, Anderson et al. analysed data on over 13 million people who had received CPR training across all 3143 counties in the USA between 2010 and 201. They concluded that the rates of CPR training were lower in black and minority ethnic communities and in lower income households.¹⁹ However, it is still unclear why people living in deprived communities are less likely to give life-saving CPR and what factors could improve this.

socioeconomically deprived circumstances to engage with bystander CPR. We defined the term 'engaging with bystander CPR' as encompassing the willingness/confidence to learn CPR, to perform bystander CPR, and to teach/ of encourage others to learn/engage with bystander CPR. METHODS This systematic review was conducted according to a registered (PBOSPEBO

according to a registered (PROSPERO CRD42017081944) and published protocol.²⁰ Studies were eligible for inclusion if they addressed any barriers or facilitators to performing bystander CPR for OHCA or being trained in CPR or training others, from a potential OHCA bystander's perspective. The following data items were sought: authors; country; year of data collection; study aims; sample; design; and barriers/ facilitators outcome measures and analysis themes. An inductive approach was used to identify barriers and facilitators in keeping with definitions proposed by Bach-Mortensen et al.21 Facilitators were defined as any factor that contributes to the delivery of bystander CPR. Barriers were defined as any factor that obstructs the delivery of bystander CPR. Studies had to either be set in a deprived area or examine a deprived population. Selected studies were limited to those published between January 2000 and December 2017 and reported on primary research. No language limitations were applied.

Search strategy and selection criteria

Searches were run on 5 December 2017 in the following: CINAHL, MEDLINE, PsycINFO, PubMed (Ahead of Print Citations and articles published in the last 6 months only) and Web of Science Core Collection (Science, Social Sciences and Arts and Humanities Citation Indices; Science and Social Science & Humanities Conference Proceedings and Books Citations Indices; and Emerging Sources Citation Index). (See sample search strategy, S1 in Supplementary Materials.) Search results were examined for relevant studies already known to the review team. Selected search terms were used in Google.co.uk, experts contacted, and reference lists checked for further studies.

We excluded studies that only addressed bystanders' use of defibrillation as our focus was on the second stage in the chain of survival ('early CPR')²¹ and the defibrillation evidence was recently reviewed.²³ We included any age group from all community settings (e.g. homes, schools, workplaces, public and private communal spaces for lay perspectives). We excluded studies with people trained and certified in CPR as part of their professional (including medical and pharmacy students), statutory (e.g. carehome staff) or voluntary roles, and studies set in medical and primary, secondary, and tertiary healthcare settings. Qualitative, quantitative, and mixed methods studies were eligible provided they collected primary data or conducted secondary analysis of existing data. Conference abstracts and unpublished 'grey' literature were also eligible and were sought through web searches, informal stakeholder interviews (from the wider project team), and our study advisory group. The reference lists of all relevant reviews were checked for additional studies to reduce the risk of bias through omitted inclusion of relevant papers.

Systematic reviews, evidence-based guidelines, and opinion pieces were excluded. Studies were eligible if they used any indicator of socioeconomic deprivation for the setting or sample including, but not limited to: educational status; employment status; income; occupation; poverty; social change; social class; social condition; or neighbourhood/area status. Study findings could report entirely from a socioeconomically deprived population or area or be segmented by a socioeconomic indicator and report from a deprived sub-population or area. For studies where the whole sample was from a socioeconomically deprived population or area, the deprivation criteria was required to be reported by

the study's authors. The review was part of a wider study to inform the development of an intervention for deprived UK communities, thus only studies from Organisation for Economic Co-operation and Development (OECD) member countries were eligible (http:// www.oecd.org/about/ membersandpartners). These tend to have high-income economics but may also have socio economic inequalities and deprived communities.

10% of records were double-screened on title and abstract by two reviewers to pilot the screening checklist. Following refinements, a further two batches of 10% of records were triple-screened on title and abstract by reviewers until high agreement was reached. Disagreements over inclusion were resolved by discussion. The remainder were singlescreened by the same three reviewers. Coding conflicts mostly related to the reason for excluding a record, not whether it should be included or excluded. A disproportionately large number of full-text studies required assessment against the inclusion criteria as it was unclear from study records whether findings were segmented by socioeconomic indicators or not. A fulltext fast-screening stage was developed to assess the deprivation inclusion criterion only. Two reviewers singlescreened full texts, using socioeconomic terms and deprivation terms from our search strategy. If no deprivation terms were identified, they closely read the sample and setting sections, and the results text and tables. The remaining full texts were assessed independently against the complete inclusion checklist by one reviewer performing the assessment and another checking the decision. Disagreements over inclusion were resolved by discussion.

Critical appraisal

A quality assessment was made of all studies, using an appropriate tool by study design: the CASP Qualitative Checklist,²⁴ the NIH Quality Assessment Tool for observational cohort and cross-sectional studies,²⁵ and the NIH Quality Assessment Tool for before-after (pre-post) studies with no control group.²⁶ We deviated from the tools stated in the protocol as none were suitable for the latter two study designs. One reviewer applied the criteria to assess the quality of the included papers. A second reviewer made a detailed check of all the assessments. Any discrepancies were resolved through consensus.

Data analysis

Relevant data were extracted into a customised form: aim, design, country and year of data collection, sample and setting, outcome measures, and findings. For studies where the sample was segmented by a socioeconomic indicator, only data related to the deprived sub-population or area were extracted. Data were extracted by one reviewer and a proportion checked for accuracy by a second. Review authors were not contacted for missing data. The extracted data (outcome measures and results) were heterogeneous, so neither a quantitative nor a qualitative metaanalysis was appropriate. To inform the narrative synthesis, data were organised thematically, and the analysis was refined through discussion. Significance and p-values are reported if they were provided in the original articles.

RESULTS

The searches produced 1219 unique records to be screened against our criteria, with 338 assessed as full-text articles (see Figure 1). Nineteen studies, reported in 21 papers, met the criteria and were included for analysis.7,27-46 Characteristics of included studies are listed in Table 1. Four studies were rated on the quality of their methods as 'good' (their findings had a low risk of bias); eight were rated as 'fair'; and seven were rated as 'poor' (their findings had a high risk of bias) (See S2 in Supplementary Materials). Most studies were conducted in Europe (n=8; from Denmark, England, France, Ireland, Italy, Spain (n=2), Sweden), followed by North America (n=6; United States), Asia (n=3; fromJapan (n=2), South Korea), and Australia (n=2). All were published articles (no grey literature), published between 2000 and 2017 with data collected between 1998 and 2015 (unreported by three studies).

Socio-demographic indicators

Six studies (reported in eight articles) specifically targeted a socioeconomically deprived population or area.^{29,36–39,41-43} Four were studies in deprived urban neighbourhoods in the USA^{29,39-43} described as having 'economic disadvantage', 'lack of health insurance', and being 'underserved': three in predominantly African American neighbourhoods (one qualitative study,^{42,43} and two surveys)^{29,39} and one qualitative study in a Latino neighbourhood.⁴¹ A UK intervention study recruited from a drug dependence unit, hostels for homeless people and primary care facilities³⁸ and a Danish intervention study took place in a rural area with lower educational levels and higher unemployment and absence due to illness rates than the rest of country^{36,37} (See S3 in Supplementary Materials for full descriptions of indicators). The remaining 13 studies were cross-sectional general population surveys and were eligible because the sample was segmented by a socio economic indicator with findings reporting potential barriers or facilitators for the lowest category of the sociodemographic indicator. Eight studies reported findings by education level,^{28,30,33,34,44–46} six by employment status or social class, 27, 28, 31, 32, 33, 40 four by household income, 30,40,45,46 and one by neighbourhood income.35 (Some studies reported by more than one indicator.)

A SYNTHESIS OF FINDINGS FROM THE INCLUDED STUDIES Learning bystander CPR

Six studies examined willingness to learn CPR, the feasibility of teaching it in schools, or barriers to learning.^{27,29,35,39,41–43} Three surveys^{27,29,39} explored willingness to learn CPR, among untrained populations. A Swedish population survey²⁷ indicated that people with a lower social-economic classification (unemployed or manual work) were willing to learn CPR, although at lower levels than professionals and students/military. Two small surveys^{29,39} set in low-income, predominantly African American urban neighbourhoods found



mixed results: a survey in Pennsylvania of a mixed age group found a high proportion of participants were willing to learn CPR;³⁹ however, results were much lower in a survey with older respondents in Florida.²⁹ The latter also found that few respondents were aware of places where CPR training took place or who to contact to attend. A further survey found that schools sited in below-average income Barcelonan neighbourhoods were as likely to be assessed as suitable for CPR training as average and aboveaverage neighbourhoods.³⁵

Two qualitative studies in lower income predominantly Latino⁴¹ and African

American^{42,43} city communities found that residents may be less motivated to participate in CPR classes when learning CPR was not a job requirement. These studies also found financial barriers related to the high cost of attending classes, and difficulties finding transport and childcare to attend.

Willingness to perform bystander CPR

Seven surveys^{31,32,34,39,45,46} and one intervention study^{22,24} measured willingness to perform bystander CPR, some for CPR on specific victims.^{36,37,39,46} Four cross-sectional surveys measured willingness to perform bystander CPR and found few differences by socio economic status. A general population survey revealed that willingness to perform CPR did not vary significantly by social class in Ireland.³² However, the survey question was asked only of those who had received CPR training in the last 5 years and respondents from lower socioeconomic classes were significantly less likely to have had training (p < 0.0001), meaning few would have been asked this. Using two hypothetical scenarios (CPR under one's own initiative and telephone-

· ·
0

Characteristics and qu	uality appraisal grading of the included stud	lies.		
Authors Country Year of data collection Quality Grading	Study's aim(s)	Sample	Design Deprivation/socio economic indicator	Barriers/facilitators outcome measures (quantitative studies) and analysis themes (qualitative studies)
Axelsson et al. ²⁷ Sweden 2000 Fair	To determine the number of CPR-trained adults in Sweden, and willingness of the non- trained to attend a CPR course. Also aimed to investigate differences (sex, age, residential area, socio economic classification and country of origin)	General population; 18–79years N = 605	Quantitative: Cross-sectional survey (postal) Analyses outcome measure by respondents' socio-demographic characteristics: Socioeconomic class	Those with no CPR training were asked if they would be willing to attend such a course
Ballesteros-Peña et al. ²⁸ Spain (Basque Country) 2015 Fair	Aimed to explore the training received, self- perceived capacitation, general knowledge and the perceptions and attitudes of the population of the Basque Country with regard to the use of CPR and AED	General population; 15–64years N = 1748	Quantitative: Cross-sectional survey (face-to- face) Analyses outcome measure by respondents' socio-demographic characteristics: Educational level and Employment status	Self-perceived capability to apply CPR procedure
Demirovic ²⁹ USA (Miami-Dade County, Florida) 1999–2000 Poor	Aimed to assess knowledge and attitudes towards CPR and AED training and use in a population sample of middle-aged and elderly African American men and women living in a large metropolitan area	US-born African American men and women, aged 50–79 years; from six areas in African American neighbourhoods of a large metropolitan region. <i>N</i> =425	Quantitative: Cross-sectional survey (face-to- face) <i>Neighbourhood (of study sample</i>): 1990 census data specific ethnic neighbourhoods; African American population	Those with no CPR training were asked if they would be willing to attend such a course Opinions and suggestions of how CPR (and related educational programmes) should be organised to make them more effective
Dwyer ³⁰ Australia (Central Queensland) 2005 Poor	Aimed to better understand barriers to family- initiated CPR and to determine whether socio-demographic variables are associated with confidence to initiate CPR on a family member	General population 18 and older <i>N</i> =1208	Quantitative: Cross-sectional survey (telephone) Analyses outcome measure by respondents' socio-demographic characteristics: <i>Educational level and Annual household</i> <i>income</i>	Confidence to initiate CPR if a family member collapses
Enami et al. ³¹ Japan (Ishikawa prefecture) 2007–2009 Fair	Aimed to examine whether ageing influences the attitudes of the population towards performing basic life support and whether the beneficial effects of exposure to basic life support courses differ between elderly and non-elderly citizens	General population driving licence applicants and drivers <i>N</i> = 22,692	Quantitative: Cross-sectional survey (paper) after course Analyses outcome measure by respondents' socio-demographic characteristics: <i>Employment status</i>	Willingness to perform BLS in two hypothetical scenarios: CPR under their own initiative and telephone-assisted compression-only CPR
Jennings et al. ³² Ireland 2008 Fair	Aimed to assess the readiness in the Irish population to (1) contact emergency services and (2) initiate CPR. Studied the proportion of the population that had attended a CPR training course in the past 5 years	General population Omnibus survey with $N = 974 \ge 15$ years. A representative sample of the population of Ireland	Quantitative: Cross-sectional survey (face-to- face) Analyses outcome measures by respondents' socio-demographic characteristics: Social class	Willingness to undertake CPR in an emergency for those trained in CPR Barriers for those trained in CPR

(Continued)

January 2023 Vol 143 No 1 | Perspectives in Public Health 47

		udies) litative	arforming arforming r person	anual or methods erceived rd, lity scores)	ners acting oncepts if	de chest de mouth- anger illing to viding sing willing ns sing willing rns fentilation	d with
		facilitators outco s (quantitative str ysis themes (qua	rted likelihood of pe r CPR rted likelihood of pe r CPR on an elderly	interference to the m d ECC techniques e combined from p eproducibility, haza ision and acceptabl	bility of schoolteachers of basic CPR contrained	n likelihood to provi isons to a stranger n likelihood to provi ventilation to a stra why they were unw why they were unw why they were unw pressions n confidence in pro npressions n confidence in pro northouth ventilation n reasons for not b e chest compressio n reasons for not b	ce in undertaking C who has overdose
		Barriers, measure and anal studies)	Self-repo bystande Self-repo bystande	Attitude/r automate (one scor efficacy, r apprehen	The feasi as provid previously	Change ii compress Change in to-mouth Reasons Provide C Change ii chest cor Change ii change ii to provide to provide to provide to provide	Confiden
		Design Deprivation/socio economic indicator	Quantitative: Cross-sectional telephone survey Analyses outcome measures by respondents' socio-demographic characteristics: <i>Educational level and Employment status</i>	Quantitative: Cross-sectional survey (online) after viewing two videos Analyses outcome measure by respondents' socio-demographic characteristics: <i>Educational level</i>	Quantitative: Cross-sectional survey (postal) Analyses outcome measure by respondents' socio-demographic characteristics: <i>average</i> income of school's neighbourhood	Quantitative: Pre- and post-intervention surveys (telephone) Educational level, Employment status and Household income (of study sample)	Quantitative: Pre- and post-intervention surveys Recruitment setting/Neighbourhood (of study sample)
	ies.	Sample	General population N = 4490 (General health survey)	General population N=1769	Head teachers of 227 public and private secondary schools	Bornholm island population N= 824 (pre) N= 815 (post) ≥15 years and over	Injecting drug users (some being current substance misuse service clients) N=107
	ality appraisal grading of the included stud	Study's aim(s)	Aimed to assess the attitudes of the public to performing CPR, perceived barriers and facilitators to performing CPR, and whether their socio-demographic and health are associated with willingness to perform bystander CPR	Aimed to test the hypothesis that the general public might perform external chest compressions (ECC) more often if they had an automated device at their disposal rather than manual ECC	Explored school interest in including a basic CPR programme in their curriculum	Nielsen et al. (2013) aimed to identify specific barriers preventing bystander interventions. Aimed to assess the effects of a 1-year targeted media campaign and education on willingness to provide basic life support and deploy an AED Nielsen et al. (2012) aimed to evaluate the effect of a multifaceted community-based intervention on bystander basic life support rates and survival to hospital discharge after OHCA of presumed cardiac aetiology	This intervention trained injecting drug users in recognising and responding to heroin overdoses (and included on-site instruction in CPR). Participants completed pre- and post- group questionnaires
Table 1 (Continued)	Characteristics and qu	Authors Country Year of data collection Quality Grading	Johnston et al. ³³ Australia (Queensland) 1998 Poor	Lapostolle et al. ³⁴ France 2010–2011 Poor	Miro et al. ³⁵ Spain (Barcelona) 2003–2004 Poor	Nielsen et al ³⁶ and Nielsen et al ³⁷ Denmark (Bornholm) 2008–2009 Fair	Phillips et al. ³⁸ UK, England (North London) Year of data collection not reported Fair

(Continued)

7	7
Q	
Ω.	
7	
)
_	
Q	
-	
<u>(</u>	

Characteristics and q	uality appraisal grading of the included stuc	lies.		
Authors Country Year of data collection Quality Grading	Study's aim(s)	Sample	Design Deprivation/socio economic indicator	Barriers/facilitators outcome measures (quantitative studies) and analysis themes (qualitative studies)
Ross et al. ³⁹ USA (Pittsburgh, Pennsylvania) Year of data collection not reported	Aimed to determine the number of individuals who have been trained in CPR and to further investigate barriers to the provision of CPR on selected victims, especially in minority communities	Two urban neighbourhoods with a high proportion of African American residents 5462 surveys mailed; N=462 surveys returned	Quantitative: Cross-sectional survey (postal) Neighbourhood (of study sample)	Those with no CPR training were asked if they would be willing to learn CPR Willingness to perform CPR Potential barriers to the performance of bystander CPR
Sasaki et al. ⁴⁰ Japan 2012 Poor	Aimed to explore factors associated with layperson confidence in performing CPR for OHCA patients, and potential emotional distress related to performing CPR	General population N = 4853	Quantitative: Cross-sectional survey (online) Analyses outcome measures by respondents' socio-demographic characteristics: <i>Employment status; Annual household</i> <i>income</i>	Whether the respondent felt confident in performing chest compressions Whether the respondent felt confident in performing mouth-to-mouth rescue breathing Potential emotional distress if a resuscitation attempt were to prove unsuccessful
Sasson et al ⁴¹ USA (Denver, Colorado) 2012 Good	The study aimed to understand common barriers that may decrease use of 911 and learning or performing bystander CPR for OHCA	Primarily Latino high-risk neighbourhoods (high OHCA incidence and low prevalence of bystander CPR) 13 years and older <i>N</i> =64	Qualitative: 6 focus groups (55 participants) and 9 key informant interviews Spatial analytical approaches were used to identify high-risk neighbourhoods Neighbourhood, Educational level and Household income (of study sample)	Analysis themes: Barriers to learning CPR, barriers to performing CPR, suggested facilitators to increase CPR training and to increase performance of bystander CPR
Sasson et al.; ⁴² King et al. ⁴³ USA (Columbus, Ohio) 2011 Good	To explore the barriers and facilitators to learning and performing CPR in three lower income, high-risk neighbourhoods To identify integral components for the design and implementation of a community-based CPR training programme for high-risk neighbourhood with lower socio economic group residents	High-risk neighbourhoods (high OHCA incidence and low prevalence of bystander CPR) <i>N</i> = 42	Qualitative: 6 focus groups (42 participants) Spatial analytical approaches were used to identify high-risk neighbourhoods (high OHCA incidence and low prevalence of bystander CPR). Five neighbourhoods were deemed high risk, and three target areas were identified for focus group interviews <i>Neighbourhood, Educational level and</i> <i>Household income (of study sample)</i>	Analysis themes: barriers to learning CPR, barriers to performing CPR, possible facilitators to learning and teaching CPR, and factors participants considered integral to the design and implementation of a successful community-based CPR training programme
Savastano and Vanni ⁷ Italy 2008–2009 Fair	To quantify the percentage of people who, having completed a basic life-saving course, would really perform CPR on an unknown adult or child. Also to evaluate the main fears, which limit performing CPR on adults or children	Lay rescuers (firefighters, flight attendants and people from the community) who had just completed 'CPR for Family and Friends' course <i>N</i> = 1000	Quantitative: Cross-sectional survey (paper) after course Analyses outcome measures by respondents' socio-demographic characteristics: <i>Educational level</i>	Willingness to perform CPR on adult patients (unknown to them) Willingness to perform CPR on paediatric patients (unknown to them)
				(Continued)

		iers/facilitators outcome isures (quantitative studies) analysis themes (qualitative lies)	fidence in their ability to perforr i in an emergency	igness to perform CPR	igness to perform hands-only stranger	
		Design Deprivation/socio economic indicator mea and stud	Quantitative: Cross-sectional survey Conf (telephone) Analyses outcome measure by respondents' socio-demographic characteristics: <i>Educational level</i>	Quantitative: Cross-sectional survey (face-to- face) Analyses outcome measure by respondents' socio-demographic characteristics: <i>Educational level</i>	Quantitative: Cross-sectional survey (face-to- Willin face) on a Analyses outcome measure by respondents' socio-demographic characteristics: <i>Educational level</i>	uut-of-hospital cardiac arrest.
	ies.	Sample	General population ≥18 years (representative sample) N = 1001	General population Quota sampling for adults aged 19 years or above living in Daegu metropolitan city <i>N</i> = 1000	Patients/visitors to an 'academic suburban emergency department' N = 532	poulmonary resuscitation: OHCA: o
uality appraisal grading of the included stud	Study's aim(s)	To define the reach of CPR training programmes in the study community and to assess residents' stated confidence in their ability to apply the skills they learned in a CPR class; also to define factors that were associated with willingness to perform CPR in an emergency situation amongst trained/ untrained participants	To determine how bystander CPR willingness is affected by factors related to CPR education (method/number of sessions, time period from the last education session) and whether AED training included	To determine current knowledge of and willingness to perform Hands-Only CPR among lay people	al defibrillator: BLS: basic life support: CPB: cardio	
	Characteristics and q	Authors Country Year of data collection Quality Grading	Sipsma et al. ⁴⁴ USA (King County, Washington) 2008 Fair	Son et al. ⁴⁵ South Korea (Daegu metropolitan city) 2012 Good	Urban et al.46 USA (Stony Brook, New York) Year of data collection not reported Poor	AED: automated externa

assisted compression-only CPR), a large survey in Japan measured willingness to perform basic life support, which it reported by occupation:³¹ there was no difference between students and those with no secure employment. The third survey, in a South Korean city, found no difference in willingness to perform CPR based on monthly income but willingness increased with level of education (from middle school graduate or under, to high school graduate, up to college graduate or higher (p < 0.001)).⁴⁵ A small survey in France found no differences by education level associated with preferred techniques to perform bystander CPR (manually, by automated device, or no preference $(p = 0.09)).^{34}$

A survey in a low-income, predominantly African American urban neighbourhood³⁹ found that a high proportion of respondents were willing to perform CPR 'on anyone'. Of the three other studies that measured willingness to perform bystander CPR on specific groups or victims, neither a lower level of income nor level of education made a difference to survey responses. A small, New York survey⁴⁶ found no significant differences in the rates of willingness to perform hands-only CPR on a stranger, when analysed by income (p = 0.82) or education level (p=0.16) Similarly, a small Italian survey on newly CPR-trained community members⁷ found no differences in rates of willingness to perform CPR on an unknown adult (p=0.670) or child (p=0.661), when analysed by educational attainment. Finally, a before-and-after study evaluated a bystander CPR intervention on an island with higher levels of unemployment and low educational attainment in relation to the rest of Denmark.^{36,37} Although the intervention had no significant effect (p = 0.15), participants were as willing to perform CPR (chest compressions only) on a stranger before as after the intervention (85% and 87%, respectively). The reasons study respondents gave for being unwilling to provide CPR to a stranger was because they did not know how to give CPR or were afraid of doing harm.³⁶

Barriers and facilitators to delivering bystander cardiopulmonary resuscitation in deprived communities: a systematic review

Щ

Confidence to perform bystander CPR

Nine studies examined respondent's confidence in administering bystander CPR.^{28,30,33,36,37,38,39,40,41,43,44} Five general population surveys reported their findings by socioeconomic status^{28,30,33,40,44} and provided a mixed picture regarding confidence to perform CPR, with some socioeconomic factors affecting confidence more than others. A general population survey in Ireland found no demographic differences, including social class, among the respondents who had been recently trained but still had concerns, like lack of confidence.33 A survey in Washington, USA,⁴⁴ found level of education had no effect on confidence to administer CPR, whether trained or not. Conversely, a small survey in Spain²⁸ found that participants with 'elementary [education] or no studies' were almost three times more likely than the rest (odds ratio (OR) = 2.7; 95% confidence interval (CI): 1.4-5.5) to respond that they felt incapable of performing CPR (90.3%) (p=0.005), but occupations were not a significant factor with regard to confidence (p = 0.05). Similarly, in a large survey in Japan,⁴⁰ working status had no effect on confidence in people's abilities (for chest compressions, p=0.178 or rescue breathing, p=0.298) but household income did; those in highest income brackets were significantly more likely than those in the lowest to be confident in their ability to perform chest compressions (p = 0.045). One survey measured confidence to initiate bystander CPR on specific people family members.³⁰ The large survey in Central Queensland, Australia, found that people with a lower education level and lower household income were less likely to feel confident at administering CPR to a family member if required. Confidence increased with income (from 63% on the lowest income to 85% on the highest) and level of education (from 57% with \leq 10 years to 77% with >15 years). Survey respondents with 11 or more vears of education were as much as two and a half times more confident to initiate CPR if a family member collapsed than those with less education (p-values unreported for individual bands, none were < 0.05).

A qualitative study^{42,43} that used focus groups with residents from an economically disadvantaged, predominantly African American community in Columbus, Ohio, indicated that low confidence in one's capability to perform bystander CPR could be caused by lack of knowledge about how and when to perform CPR and confusion caused by guidelines changing. A small survey with older respondents in a lower income, predominantly African American Florida city neighbourhood found between 16% and 20%, who had been trained in CPR did 'not feel at all comfortable with their CPR skills' (p. 66).39

Finally, two training interventions targeted at increasing engagement with CPR facilitated an improvement in their confidence and capability to perform bystander CPR.^{36,37,38} In a post-training survey in England, more injecting drug users felt confident to undertake CPR with a person who had overdosed than before³⁸ and residents on a Danish island with higher unemployment and lower educational attainment than the national average had a small increase in confidence at providing CPR after training and media campaigns.^{36,37}

Likelihood of performing bystander CPR

Two studies examined the perceived or declared likelihood of administering bystander CPR if the situation arose.33,41 In a large household survey in Queensland, Australia,³³ both employment status (p < 0.001) and education level (p < 0.001) were found to be significantly associated with the likelihood of performing bystander CPR. The article did not report the significance of individual levels of education or occupation, but proportions for declaring they were 'extremely likely' to perform bystander CPR ranged from those on home duties (44%) and unemployed (52%) to full-time workers (62%) and students (67%); and from those who had completed primary school (38%) to 'trade/tech/dipl.' (63%) education level. The same survey identified that neither employment status nor education level had a significant association with whether they were more or less likely to perform CPR on an elderly victim. In a

qualitative study⁴¹ in a low-income, predominantly Latino city neighbourhood in Colorado, a commonly cited barrier affecting residents' likelihood of performing bystander CPR was the age and sex of the bystander being different to the victim's. Residents also expressed that the existence of Latino and Black gangs in their own communities may make bystanders less likely to get involved in performing CPR, depending on the race of the victim involved.

Other factors influencing delivery of bystander CPR

Risk to personal health and safety Five studies raised the issue of personal risks from performing bystander CPR.^{32,36,37,39,41-43} Fear of catching a disease or infection from administering mouth-to-mouth ventilation was raised by focus group participants in two qualitative studies^{41–43} and in one small survey,³⁹ all conducted in lower income, predominantly minority ethnic neighbourhoods in three US cities. Fear was also cited as a barrier to performing CPR by some survey respondents on an island with higher unemployment and lower educational attainment compared with the rest of Denmark gave for being unwilling to provide CPR to a stranger;^{36,37} and by participants in a household survey in Ireland who had been recently trained in CPR.³² Two qualitative studies^{41–43} identified that being in a location which may place the bystander in danger, for example, at risk of being attacked or robbed was also perceived as a potential barrier to performing CPR.

Fear of legal consequences

Four studies, three from the USA, revealed that fear of legal consequences may be a barrier to bystander CPR.^{32,39,41–43} Two qualitative studies reported that the fear of lawsuits and legal consequences acted as a main barrier to performing bystander CPR among the participants from lower income Latino⁴¹ and African American^{42,43} urban neighbourhoods. A small survey with older respondents in a lower income Florida city neighbourhood found the same,³⁹ as did the household

survey in Ireland, for those who had been recently trained but had concerns.³²

Lack of community connectedness

Two qualitative studies found that engaging in CPR was difficult in the economically disadvantaged, predominantly African American community in Columbus, Ohio,^{42,43} and the predominantly Latino community in Denver, Colorado,⁴¹ owing to a heightened lack of community connectedness. In focus groups in both cities, some residents conveyed that they did not know their neighbour or felt emotionally disconnected from them. This disconnection made them more reluctant to assist a victim in the street whom they did not know.

Cultural barriers for a specific community

A qualitative study,⁴¹ where participants were from lower income, predominantly Latino neighbourhoods in Denver, Colorado (USA), raised additional culturally specific factors for engaging with bystander CPR in their communities. Participants feared that if they administered bystander CPR, they would be asked for identification or blamed for the victim's condition when police and paramedics arrived on scene. Language concerns and touching someone in a way that could be perceived as inappropriate were cited as additional barriers to performing bystander CPR.

DISCUSSION

The review identified key barriers and facilitators that impact on the ability of individuals who live in socioeconomically deprived circumstances in OECD nations to engage in bystander CPR: the willingness to learn or perform CPR; the confidence to perform CPR; and selfreported likelihood of performing CPR. Additional identified barriers and facilitators to engaging with CPR appeared to be specific for individuals from socioeconomically deprived backgrounds or areas. Few studies measured the self-reported likelihood of performing CPR.

As previously identified,¹⁸ the findings indicate that many people in within the included study settings were willing to learn CPR, regardless of their level of education or income. However, lower community affluence or socioeconomic backgrounds of participants appeared to impact on the likelihood of individuals receiving training.^{19,47} Findings from some studies suggest that low levels of motivation to become trained if it is not a job requirement, unaffordability of training, or inaccessibility of training due to lack of childcare, all acted as barriers with their study contexts.^{41–43} The review identified that individuals' confidence in their ability to perform bystander CPR was mixed. Some measures of socioeconomic status appear to affect confidence levels more than others. Overall, populations with low education or household income levels appear to have low confidence in administering CPR to either a family member or a stranger. This review revealed other barriers to engaging with CPR: the risk to personal health linked to the environment, a fear of legal consequences, a lack of community cohesion, and a number of cultural barriers. Moreover, in some of the deprived communities that were studied in this review, there was a perception potential risk to bystanders' safety from helping victims in situations where they themselves could be robbed or attacked.

Since the systematic search for studies in December 2017 and the results were synthesised, two relevant studies have been published. (The search strategy was rerun in Medline only, in August 2020.) A cross-sectional survey of public gatherings in Baltimore, USA,48 found that those from the highpoverty areas (27% of all respondents) preferred instructor-led CPR training and that most disliked the idea of training occurring at a local learning station (at events or in shops) and preferred libraries. Non-college graduates (46% of all respondents) preferred school settings as well as instructor-led education. The survey found that preferences for training locations and formats were comparable across all respondents. A small prospective survey by the same authors conducted before and after free community compression-only CPR classes in Baltimore⁴⁹ found that a lower level of education (29% of all

respondents) increased the likelihood that participants feared being sued or risk of disease or hurting someone and were unlikely to perform CPR on strangers or family, even after training. Household income was not found to affect the likelihood of performing CPR. Neither of these studies substantially alter our results.

The findings from this review have a number of implications for the development of interventions aimed at improving bystander CPR in deprived communities. Although such populations show a willingness to learn CPR, they are often among the least trained in CPR. Therefore, interventions to support bystander CPR must make training more flexible, affordable, and accessible to people in deprived communities. More education needs to occur and to be targeted at deprived communities focusing not only the practical skills to give bystander CPR but also the confidence that anyone may be 'CPR ready' and capable to give bystander CPR, even if they are weaker or disabled, or at least that they may be able to give help. The findings of this review informed the analysis of a qualitative research study on bystander CPR with deprived communities in Scotland which identified similar individual and environmental barriers to be targeted through tailored interventions.50

LIMITATIONS

A key limitation of the review's method was the range of proxy sociodemographic indicators used for signifying a socially or economically disadvantaged community. Some of the included studies were unambiguously conducted with people living in deprived circumstances. Other studies employed single indicators like low educational attainment or large geographic areas where employment status and incomes were averaged across all residents are more uncertain for labelling the respondents as living in deprived circumstances. Other limitations include the potential for introducing reviewer bias and missing relevant studies by singlescreening at the 'fast screening' stage, as none of the papers excluded were double-checked. Few of the included

studies' main aims were to identify barriers and facilitators of bystander CPR in deprived communities. Many of the included (and excluded) surveys collected barriers and facilitators data but did not analyse all outcome measures by socio-demographics. Finally, thresholds for deprivation may not be defined in the same way across countries. This had an impact on the way the studies presented their results and made it difficult to compare findings.

CONCLUSION

This review suggests that people living in deprived communities in different contexts face significant and specific barriers to becoming 'CPR ready'. Tailored interventions for deprived communities are required to increase engagement with bystander CPR, including information and awareness raising (through contextspecific and community-led social marketing campaigns) and more innovative and accessible CPR training (through peer and social network interventions, online, shorter and cheaper CPR training, support for childcare, and transport costs).

ACKNOWLEDGEMENTS

Our thanks to Anne Marie MacKintosh (ISMH) for assistance with interpreting some of the studies' statistical data and to the advisory group for sharing relevant papers. We acknowledge members of the wider study for their support: Martine Stead, Douglas Eadie, and Nathan Critchlow (ISMH, University of Stirling), and Gareth Clegg, Liz Hasseld, and Lisa MacInnes (Resuscitation Research Group, University of Edinburgh).

CONFLICT OF INTEREST

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

FUNDING

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The systematic review was conducted as part of a project funded by the Chief Scientist Office HIPS /17/10. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The corresponding author had final responsibility for the decision to submit for publication.

ORCID ID

I Uny D https://orcid.org/0000-0002-9548-5332

K Angus () https://orcid.org/0000-0002-5351-4422

SUPPLEMENTAL MATERIAL

Supplemental material for this article is available online.

References

- Myat A, Song K-J, Rea T. Out-of-hospital cardiac arrest: current concepts. *The Lancet* 2018;**391**:970–9.
- Berdowski J, Berg RA, Tijssen JGP et al. Global incidences of out-of-hospital cardiac arrest and survival rates: systematic review of 67 prospective studies. *Resuscitation* 2010;81(11):1479–87.
- Sasson C, Rogers MA, Dahl J et al. Predictors of survival from out-of-hospital cardiac arrest: a systematic review and meta-analysis. *Circ Cardiovasc Qual Outcomes* 2010;**3**(1):63–81.
- Coons SJ, Guy MC. Performing bystander CPR for sudden cardiac arrest: behavioral intentions among the general adult population in Arizona. *Resuscitation* 2009;80(3):334–40.
- Axelsson A, Thoren A, Holmberg S et al. Attitudes of trained Swedish lay rescuers toward CPR performance in an emergency: a survey of 1012 recently trained CPR rescuers. *Resuscitation* 2000;44(1):27–36.
- Schmid KM, Mould-Millman NK, Hammes A et al. Barriers and facilitators to community CPR education in San Jose, Costa Rica. Prehospital Disaster Med 2016;31:509–15.
- Savastano S, Vanni V. Cardiopulmonary resuscitation in real life: the most frequent fears of lay rescuers. *Resuscitation* 2011;82(5):568– 71.
- Mitchell MJ, Stubbs BA, Eisenberg MS. Socioeconomic status is associated with provision of bystander cardiopulmonary resuscitation. *Prehosp Emerg Care* 2009;**13**(4):478–86.
- Sasson C, Haukoos JS, Eigel B et al. The HANDDS program: a systematic approach for addressing disparities in the provision of bystander cardiopulmonary resuscitation. Acad Emerg Med 2014;21(9):1042–9.

- Moon S, Bobrow BJ, Vadeboncoeur TF et al. Disparities in bystander CPR provision and survival from out-of-hospital cardiac arrest according to neighborhood ethnicity. Am J Emerg Med 2014;32(9):1041–5.
- Rivera NT, Kumar SL, Bhandari RK et al. Disparities in Survival with Bystander CPR following cardiopulmonary arrest based on neighborhood characteristics. *Emerg Med Int* 2016;2016:6983750.
- Hasselqvist-Ax I, Riva G, Herlitz J et al. Early cardiopulmonary resuscitation in out-ofhospital cardiac arrest. N Engl J Med 2015;372(24):2307–15.
- Stromsoe A, Svensson L, Axelsson AB et al. Improved outcome in Sweden after out-ofhospital cardiac arrest and possible association with improvements in every link in the chain of survival. *Eur Heart J* 2015;36(14):863–71.
- Wissenberg M, Lippert FK, Folke F et al. Association of national initiatives to improve cardiac arrest management with rates of bystander intervention and patient survival after out-of-hospital cardiac arrest. *JAMA* 2013;310(13):1377–84.
- Sasson C, Magid DJ, Chan P, et al. Association of neighborhood characteristics with bystander-initiated CPR. *N Engl J Med* 2012;367(17):1607–15.
- Moncur L, Ainsborough N, Ghose R et al. Does the level of socioeconomic deprivation at the location of cardiac arrest in an English region influence the likelihood of receiving bystander-initiated cardiopulmonary resuscitation? *Emerg Med J* 2016;33:105–8.
- Blewer AL, Ibrahim SA, Leary M et al. Cardiopulmonary resuscitation training disparities in the United States. J Am Heart Assoc 2017;6:14.

- Dobbie F, MacKintosh AM, Clegg G et al. Attitudes towards bystander cardiopulmonary resuscitation: results from a cross-sectional general population survey. *PLoS ONE* 2018;**13**(3):e0193391.
- Anderson ML, Cox M, Al-Khatib SM et al. Rates of cardiopulmonary resuscitation training in the United States. *JAMA Intern Med* 2014;174(2):194-201.
- Dobbie F, Angus K, Uny I *et al.* Protocol for a systematic review to identify the barriers and facilitators to deliver bystander cardiopulmonary resuscitation (CPR) in disadvantaged communities. *Syst Rev* 2018;**7**:143.
- Bach-Mortensen AM, Lange BCL, Montgomery P. Barriers and facilitators to implementing evidence-based interventions among third sector organisations: a systematic review. *Implement Sci* 2018;13:103.
- Nolan J, Soar J, Eikeland H. The chain of survival. *Resuscitation* 2006;**71**:270–1.
- Smith CM, Lim Choi Keung SN, Khan MO et al. Barriers and facilitators to public access defibrillation in out-of-hospital cardiac arrest: a systematic review. Eur Heart J Qual Care Clin Out 2017;3:264–73.
- Critical Appraisal Skills Programme. CASP Qualitative Checklist, 2018. Available online at: https://casp-uk.net/wp-content/uploads/2018/ 01/CASP-Qualitative-Checklist-2018.pdf (last accessed 12 July 2021).
- National Institutes of Health. Quality assessment tool for observational cohort and cross-sectional studies. Bethesda, MD: National Institutes of Health, US Department of Health & Human Services, 2014. Available online at: https://www.nhlbi.nih.gov/health-pro/ guidelines/in-develop/cardiovascular-riskreduction/tools (last accessed 12 July 2021).

- National Institutes of Health. Quality assessment tool for before-after (pre-post) studies with no control group. Bethesda, MD: National Institutes of Health, US Department of Health & Human Services, 2014. Available online at: https://www.nhlbi.nih.gov/health-pro/ guidelines/in-develop/cardiovascular-riskreduction/tools (last accessed 12 July 2021).
- Axelsson AB, Herlitz J, Holmberg S et al. A nationwide survey of CPR training in Sweden: foreign born and unemployed are not reached by training programmes. *Resuscitation* 2006;70:90–7.
- Ballesteros-Pena S, Fernandez-Aedo I, Perez-Urdiales I et al. Knowledge and attitudes of citizens in the Basque Country (Spain) towards cardiopulmonary resuscitation and automatic external defibrillators. *Med Intensiva* 2016;40(2):75–83.
- Demirovic J. Cardiopulmonary resuscitation programs revisited: results of a community study among older African Americans. Am J Geriatr Cardiol 2004;13(4):182–7.
- Dwyer T. Psychological factors inhibit family members' confidence to initiate CPR. *Prehosp Emerg Care* 2008;12(2):157–61.
- Enami M, Takei Y, Inaba H et al. Differential effects of ageing and BLS training experience on attitude towards basic life support. *Resuscitation* 2011;82(5):577–83.
- Jennings S, Hara TO, Cavanagh B et al. A national survey of prevalence of cardiopulmonary resuscitation training and knowledge of the emergency number in Ireland. *Resuscitation* 2009;80(9):1039–42.
- Johnston TC, Clark MJ, Dingle GA et al. Factors influencing Queenslanders' willingness to perform bystander cardiopulmonary resuscitation. *Resuscitation* 2003;56(1):67–75.

- Lapostolle F, Bertrand P, Agostinucci JM et al. Web-based general public opinion study of automated versus manual external chest compression. *Emerg Med J* 2014;31(6):488–92.
- Miro O, Jimenez-Fabrega X, Espigol G et al. Teaching basic life support to 12-16 year olds in Barcelona schools: Views of head teachers. *Resuscitation* 2006;70(1):107–16.
- Nielsen AM, Isbye DL, Lippert FK et al. Can mass education and a television campaign change the attitudes towards cardiopulmonary resuscitation in a rural community. ? Scand J Trauma Resusc Emerg Med 2013;21:8.
- Nielsen AM, Lou Isbye D, Knudsen Lippert F et al. Engaging a whole community in resuscitation. *Resuscitation* 2012;83(9):1067–71.
- Phillips P, Glover C, Allan T et al. Using a group approach to preventing heroin overdose in North London. *Drugs: Educ, Prev Pol* 2009;16:328–42.
- Ross CA, Winter MK, Mossesso VN. Bystander CPR in two predominantly African American communities. *Topic Emerg Med* 2000;22:63–8.
- Sasaki M, Ishikawa H, Kiuchi T et al. Factors affecting layperson confidence in performing resuscitation of out-ofhospital cardiac arrest patients in Japan. *Acute Med Surg* 2015;2:183–9.
- Sasson C, Haukoos JS, Ben-Youssef L et al. Barriers to calling 911 and learning and performing cardiopulmonary resuscitation for residents of primarily Latino, high-risk neighborhoods in Denver, Colorado. Ann Emerg Med 2015;65(5):545–52.
- 42. Sasson C, Haukoos JS, Bond C et al. Barriers and facilitators to learning and performing cardiopulmonary resuscitation in neighborhoods with low bystander

cardiopulmonary resuscitation prevalence and high rates of cardiac arrest in Columbus, OH. *Circ: Cardiovasc Qual Out* 2013;6:550–8.

- King R, Heisler M, Sayre MR e t a l. Identification of factors integral to designing community-based CPR interventions for highrisk neighborhood residents'. *Prehospital Emergency Care* 2015;19:308–12.
- Sipsma K, Stubbs BA, Plorde M. Training rates and willingness to perform CPR in King County, Washington: a community survey. *Resuscitation* 2011;82(5):564–7.
- Son JW, Ryoo HW, Moon S et al. Association between public cardiopulmonary resuscitation education and the willingness to perform bystander cardiopulmonary resuscitation: a metropolitan citywide survey. *Clin Exp Emerg Med* 2017;4(2):80–7.
- Urban J, Thode H, Stapleton E et al. Current knowledge of and willingness to perform Hands-Only (TM) CPR in laypersons. *Resuscitation* 2013;84: 1574–8.
- Hawkes CA, Brown TP, Booth S *et al.* Attitudes to cardiopulmonary resuscitation and defibrillator use: a survey of UK adults in 2017. *J Am Heart Assoc* 2019;**8**:e008267.
- Fratta KA, Bouland AJ, Vesselinov R et al. Evaluating barriers to community CPR education. Am J Emerg Med 2020;38(3): 603–9.
- Fratta KA, Bouland AJ, Lawner BJ et al. Barriers to bystander CPR: evaluating socioeconomic and cultural factors influencing students attending community CPR training. Am J Emerg Med 2019;37(1):159–61.
- Dobbie F, Uny I, Eadie D *et al.* Barriers to bystander CPR in deprived communities: findings from a qualitative study. *PLoS ONE* 2020;**15**:e0233675.