

LAMPIRAN

Lampiran 1. Surat Persetujuan Menjadi Responden

SURAT PERSETUJUAN

(INFORMED CONSENT)

Saya yang bertanda tangan di bawah ini,

Nama :
Usia :
Alamat :

Menyatakan bahwa :

1. Saya telah mendapatkan penjelasan mengenai tujuan dan manfaat penelitian dari :

Nama : Amaliya Ilma Nafia
NIM : P17110173048
Jurusan : D-III Gizi Politeknik Kesehatan Kemenkes Malang
Judul : Kebiasaan Mengonsumsi Kopi dan Merokok sebagai Faktor Eksternal Penyebab Hipertensi

2. Setelah saya memahami penjelasan tersebut, dengan penuh kesadaran dan tanpa paksaan dari siapapun, saya bersedia ikut serta dalam penelitian ini dengan kesepakatan sebagai berikut :
 - a. Data yang diperoleh dari penelitian ini akan dijaga kerahasiaannya dan hanya dipergunakan untuk kepentingan ilmiah.
 - b. Apabila suatu waktu saya merasa dirugikan dalam bentuk apapun, saya berhak membatalkan persetujuan ini serta berhak untuk mengundurkan diri.

Demikian surat persetujuan ini saya buat dengan sesungguhnya dan dapat digunakan sebagaimana mestinya.

Malang,

Responden

(.....)

Lampiran 2. Data hasil penelitian

No.	Nama	Usia	Riwayat hipertensi keluarga	Status gizi	Tekanan Darah	Konsumsi kopi per hari	Merokok per hari	Recall					
								E	P	L	KH	Vit C	Na
1.	Tn. MY	30 tahun	Tidak	Obesitas	148/90 mmHg	4 gelas	10-20 batang	1855.4	66	69.05	243.25	45.5	681.25
2.	Tn. S	44 tahun	Ya	Normal	126/80 mmHg	1 gelas	<10 batang	2086.5	61.35	64.65	321.6	41.55	455.05
3.	Tn. A	30 tahun	Tidak	Kurus	122/80 mmHg	3 gelas	<10 batang	1318.95	49.4	26.8	216.65	21.3	322.65
4.	Tn. IS	40 tahun	Ya	Obesitas	140/90 mmHg	3 gelas	10-20 batang	2112.2	57.35	49.3	356.1	21.6	763.85
5.	Tn. DP	39 tahun	Tidak	Normal	120/90 mmHg	5 gelas	10-20 batang	1530.8	51.4	34.15	250.1	7.55	657.05

Lampiran 3. Hasil Uji Statistika

WEIGHT BY Frekuensi.

CROSSTABS

/TABLES=Kopi BY Hipertensi

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Output Created		08-JUN-2020 06:41:09
Comments		
Input	Active Dataset	DataSet2
	Filter	<none>
	Weight	Frekuensi
	Split File	<none>
	N of Rows in Working Data File	3
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.

Syntax	CROSSTABS	
	/TABLES=Kopi BY Hipertensi	
	/FORMAT=AVALUE TABLES	
	/STATISTICS=CHISQ	
	/CELLS=COUNT	
	/COUNT ROUND CELL.	
Resources	Processor Time	00:00:00.02
	Elapsed Time	00:00:00.01
	Dimensions Requested	2
	Cells Available	131029

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Kopi * Hipertensi	5	100.0%	0	0.0%	5	100.0%

Kopi * Hipertensi Crosstabulation

Count

		Hipertensi		Total
		Prahipertensi	Hipertensi Stadium 1	
Kopi	1-2 gelas	1	0	1
	3-5 gelas	2	2	4
Total		3	2	5

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.833 ^a	1	.361		
Continuity Correction ^b	.000	1	1.000		
Likelihood Ratio	1.185	1	.276		
Fisher's Exact Test				1.000	.600
Linear-by-Linear Association	.667	1	.414		
N of Valid Cases	5				

a. 4 cells (100.0%) have expected count less than 5. The minimum expected count is .40.

b. Computed only for a 2x2 table

WEIGHT BY Frekuensi.

CROSSTABS

/TABLES=Rokok BY Hipertensi

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Output Created		08-JUN-2020 06:36:49
Comments		
Input	Active Dataset	DataSet1
	Filter	<none>
	Weight	Frekuensi
	Split File	<none>
	N of Rows in Working Data File	3
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.

Syntax	CROSSTABS	
	/TABLES=Rokok BY Hipertensi	
	/FORMAT=AVALUE TABLES	
	/STATISTICS=CHISQ	
	/CELLS=COUNT	
	/COUNT ROUND CELL.	
Resources	Processor Time	00:00:00.02
	Elapsed Time	00:00:00.01
	Dimensions Requested	2
	Cells Available	131029

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Rokok * Hipertensi	5	100.0%	0	0.0%	5	100.0%

Rokok * Hipertensi Crosstabulation

Count

		Hipertensi		Total
		Prahipertensi	Hipertensi Stadium 1	
Rokok	<10 batang per hari	2	0	2
	10-20 batang per hari	1	2	3
Total		3	2	5

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	2.222 ^a	1	.136		
Continuity Correction ^b	.313	1	.576		
Likelihood Ratio	2.911	1	.088		
Fisher's Exact Test				.400	.300
Linear-by-Linear Association	1.778	1	.182		
N of Valid Cases	5				

a. 4 cells (100.0%) have expected count less than 5. The minimum expected count is .80.

b. Computed only for a 2x2 table

CROSSTABS

/TABLES=Protein BY Hipertensi

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Output Created		26-JUN-2020 07:43:13
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	Frekuensi
	Split File	<none>
	N of Rows in Working Data File	4
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.

Syntax	CROSSTABS	
	/TABLES=Protein BY Hipertensi	
	/FORMAT=AVALUE TABLES	
	/STATISTICS=CHISQ	
	/CELLS=COUNT	
	/COUNT ROUND CELL.	
Resources	Processor Time	00:00:00.02
	Elapsed Time	00:00:00.21
	Dimensions Requested	2
	Cells Available	131029

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Asupan protein * Tekanan darah	5	100.0%	0	0.0%	5	100.0%

Asupan protein * Tekanan darah Crosstabulation

Count

		Tekanan darah		Total
		Prahipertensi	Hipertensi stadium 1	
Asupan protein	Normal	1	1	2
	Defisit tingkat sedang	2	1	3
Total		3	2	5

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.139 ^a	1	.709		
Continuity Correction ^b	.000	1	1.000		
Likelihood Ratio	.138	1	.710		
Fisher's Exact Test				1.000	.700
Linear-by-Linear Association	.111	1	.739		
N of Valid Cases	5				

a. 4 cells (100.0%) have expected count less than 5. The minimum expected count is .80.

b. Computed only for a 2x2 table

WEIGHT BY Frekuensi.

CROSSTABS

/TABLES=Lemak BY Hipertensi

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ

/CELLS=COUNT

/COUNT ROUND CELL

Crosstabs

Notes

Output Created		26-JUN-2020 07:45:51
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	Frekuensi
	Split File	<none>
	N of Rows in Working Data File	3
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.

Syntax	CROSSTABS	
	/TABLES=Lemak BY Hipertensi	
	/FORMAT=AVALUE TABLES	
	/STATISTICS=CHISQ	
	/CELLS=COUNT	
	/COUNT ROUND CELL.	
Resources	Processor Time	00:00:00.03
	Elapsed Time	00:00:00.12
	Dimensions Requested	2
	Cells Available	131029

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Asupan lemak * Tekanan darah	5	100.0%	0	0.0%	5	100.0%

Asupan lemak * Tekanan darah Crosstabulation

Count

		Tekanan darah		Total
		Prahipertensi	Hipertensi stadium 1	
Asupan lemak	Normal	0	1	1
	Defisit tingkat berat	3	1	4

Total	3	2	5
-------	---	---	---

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.875 ^a	1	.171		
Continuity Correction ^b	.052	1	.819		
Likelihood Ratio	2.231	1	.135		
Fisher's Exact Test				.400	.400
Linear-by-Linear Association	1.500	1	.221		
N of Valid Cases	5				

a. 4 cells (100.0%) have expected count less than 5. The minimum expected count is .40.

b. Computed only for a 2x2 table